

Synthesis and SAR of b-Annulated 1,4-Dihydropyridines Define Cardiomyogenic Compounds as Novel Inhibitors of TGFbeta Signaling.

Journal: J Med Chem

Publication Year: 2012

Authors: Dennis Schade, Marion Lanier, Erik Willems, Karl Okolotowicz, Paul Bushway, Christine Wahlquist, Cynthia Gilley, Mark Mercola, John R Cashman

PubMed link: 23130626

Funding Grants: Chemical Genetic Approach to Production of hESC-derived Cardiomyocytes, Burnham Institute
CIRM Stem Cell Training Grant (Type II)

Public Summary:

We used high throughput screening of chemical library to discover small drug-like molecules that promote progenitor cell to cardiomyocyte differentiation. One active compound is described in this paper. The molecule, named ITD-1, is the first selective inhibitor of Transforming Growth Factor-beta signaling. Other inhibitors of TGF-beta act by blocking the catalytic activity of its cell surface receptor, and do not distinguish between it and closely related receptors. ITD-1, in contrast, selectively causes the type II TGF-beta receptor (TGFB2) for internalization and degradation. Inhibition of TGF-beta constitutes a critical step necessary for the differentiation of uncommitted progenitor cells into cardiomyocytes. Furthermore, ITD-1 might be developed further into an drug for blocking TGF-beta function in vivo in a number of clinical settings.

Scientific Abstract:

A medium-throughput murine embryonic stem cell (mESC)-based high-content screening of 17000 small molecules for cardiogenesis led to the identification of a b-annulated 1,4-dihydropyridine (1,4-DHP) that inhibited transforming growth factor beta (TGFbeta)/Smad signaling by clearing the type II TGFbeta receptor from the cell surface. Because this is an unprecedented mechanism of action, we explored the series' structure-activity relationship (SAR) based on TGFbeta inhibition, and evaluated SAR aspects for cell-surface clearance of TGFbeta receptor II (TGFB2) and for biological activity in mESCs. We determined a pharmacophore and generated 1,4-DHPs with IC(50)s for TGFbeta inhibition in the nanomolar range (e.g., compound 28, 170 nM). Stereochemical consequences of a chiral center at the 4-position was evaluated, revealing 10- to 15-fold more potent TGFbeta inhibition for the (+)- than the (-) enantiomer. This stereopreference was not observed for the low level inhibition against Activin A signaling and was reversed for effects on calcium handling in HL-1 cells.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/synthesis-and-sar-b-annulated-14-dihydropyridines-define-cardiomyogenic>